

Beef Slaughter Plant Application

Carcass after removal of hide

Post skinning wash

Inhibitor or Microbial Blocking Agent (carageenan or CMC or dextran sulfate) dispensed in a spray box

Evisceration

Splitting

Final wash
(water spray)

Removal of bacterial contamination with
arginine and Tween 80/NaCl or phosphate buffer with Tween/NaCl

Chill

Fabrication

Trimmings; Primals

Inhibitor or Microbial Blocking Agent (carageenan or CMC or dextran sulfate) dispensed as spray or immersion
(Anti-bacterial compound may also be incorporated)

Packaging

Distribution, retailing

FIG. 1

BEST AVAILABLE COPY

Poultry (Chicken) Slaughter Plant Application

Kill Room

Scalding/defeathering

Evisceration

Chlorination

(high pressure chlorinated water jets thoroughly clean the entire bird)

Inspection

(for fecal contamination before chilling)

Chilling

(The birds body temp is about 98° F and must be lowered to 40° F or below before processing. Birds are sent to soak in the chiller for 73 minutes. Each chiller holds 20,000 gals. of chlorinated water.)

Inhibitor or Microbial Blocking Agent (carageenan or CMC or dextran sulfate) dispensed in a spray box

Grading

Birds are rehung in shackles to be graded.

Packaging of Whole Birds

or

Cut-up

2nd Inhibitor or MBA (carageenan or CMC or dextran sulfate) application to cut-ups

(anti-bacterial agent such as Cetyl Pyridium Chloride may be added)
[MBA= Microbial Blocking Agent; CMC= carboxy methyl cellulose]

Boning and trimming

Packaging of cut-up parts

FIG. 2

Swine Slaughter Plant Application in Clean Room

Shaving

Carcass Washing

Head Removal, Brisket sawing, Debunging

Carcass Opening

Evisceration

Carcass Splitting

Trimming

Stamping

Final Carcass Washing
(spraying with room temp. water)

Inhibitor or Microbial Blocking Agent (carageenan or CMC or dextran sulfate) dispensed in a spray box
(also prevents dehydration)

Chilling
(dry air blast at 2 degrees, overnight)

Cutting

Inhibitor or MBA (carageenan or CMC or dextran sulfate) application
(anti-bacterial agent such as Cetyl Pyridium Chloride may be added)
[MBA= Microbial Blocking Agent; CMC= carboxy methyl cellulose]

Packaging or Processing

FIG. 3

Detachment for Sampling
[Laboratory Analysis]

**Coat tissues with Inhibitor or Microbial Blocking Agent
(Carragenan, CMC or dextran sulfate)**

**Detach bacteria
Guanidine-HCl, pH 4.8
or
Phosphate buffer-NaCl-Tween 80**

Enrich bacterial growth

Analyze bacteria using

- 1. Traditional Plate Method**
or
2. Isolate target bacteria
immunomagnetic beads
or
imuno-affinity silica gel
or
- 3. Biosensor screening (BIAcore, etc.)**
or
- 4. Other microbial screening and testing methods**

FIG. 4

**Detachment for *In Situ* Sampling in
Slaughter Plants**

**Spray a 300 cm area with Carrageenans,
or Carboxy methyl Cellulose or dextran sulfate**

**Spray same area with arginine
to loosen the bound bacteria**

**Soak sponge with
0. 05% Tween 80 (Span 80) and 1% NaCl
or
phosphate buffer with NaCl and Tween 80**

**Wipe off bacteria from treated surface
with pre-soaked sponge
or
with filter paper (e.g. cellulose or polycarbonate membranes)
or
with vacuum (with filter) to remove bacteria from carcass surface
remove vacuum filter for entrapped bacteria**

Enrich or Extract bacteria from sponge or filter or membranes


Analyze bacteria with

- 1. Traditional Plate Methods**
or
- 2. Isolate target bacteria
with immunomagnetic beads**
or
- 3. Biosensor screening (BIAcore, etc.)**
or
- 4. Other microbial screening and testing methods**

FIG. 5

SCANNED. # 14


Heparan Sulfate



-[iduronate-2 sulfate- α 1,4 N-sulfate glucosamine-6-sulfate α 1,4]-
-[glucuronate- β 1,4...]-


FIG. 6A

Kappa Carrageenan




3,6-anhydro-D-galactose D-galactose-4-sulphate

Iota Carrageenan



3,6-anhydro-D-galactose-2-Sulphate D-galactose-4-sulphate

Lambda Carrageenan



D-galactose-2,6-disulphate D-galactose-2-sulphate

FIG. 6B

FIG. 7A




FIG. 7B




FIG. 8A




FIG. 8B





FIG. 8B

FIG. 8C



09822947 - 04004

FIG. 8D





FIG. 8E



© 2003 Wiley Periodicals, Inc. *J Biostat* 2003; 29: 100–110

3M ECOLI 24E-42 PG 3M ECOLI 24E-42 PG




FIG. 9A




FIG. 9B

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- BLACK BORDERS**
- IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- FADED TEXT OR DRAWING**
- BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- SKEWED/SLANTED IMAGES**
- COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- GRAY SCALE DOCUMENTS**
- LINES OR MARKS ON ORIGINAL DOCUMENT**
- REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.